

SOME PROPERTIES CONCERNING THE HYPERSURFACES OF A WEYL SPACE

NİL KOFOĞLU*

Mimar Sinan Güzel Sanatlar Üniversitesi, Fen-Edebiyat Fakültesi, Matematik Bölümü, Beşiktaş-İSTANBUL

Geliş/Received: 21.12.2004 Kabul/Accepted: 03.10.2005

ABSTRACT

Let W_n be a hypersurface of the Weyl space W_{n+1} . Let $v_i (i = 1, 2, \dots, n)$ be tangent vector fields belonging to W_n and n be the normalized normal vector field of W_n . Consider the $(n+1)$ -net $(v_1, v_2, \dots, v_n, n)$. By using the prolonged covariant differentiation, we first obtain the set of formulas corresponding to the Frenet formulas for W_n associated with a curve C on W_n having the tangent vector field v_1 . We then, derive two invariants concerning the orthogonal ennuple $(v_1, v_2, \dots, v_n, v_i)$, $v_i (i = 1, 2, \dots, n)$ being differentiable vector fields on W_n .

Keywords: Weyl space, Net of curves, Prolonged covariant derivative.
MSC number/numarası: 57R55, 58A99.

BİR WEYL UZAYININ GÖZÖNÜNE ALINAN HİPER YÜZEYLERİNİN BAZI ÖZELLİKLERİ

ÖZET

W_n, W_{n+1} Weyl uzayının bir hiperyüzeyi olsun. $v_i (i = 1, 2, \dots, n)$, W_n 'e ait teğet vektör alanları ve n , W_n 'in normalize edilmiş normal vector alanı olsun. $(v_1, v_2, \dots, v_n, n)$ $(n+1)$ -li şebeke göz önüne alınsın. Genelleştirilmiş kovaryant türev kullanılarak, önce W_n hiperyüzeyinin bir C eğrisinin v_1 teğet vector alanına bağlı olarak Frenet formüllerine tekabül eden formüller elde edilmiştir. Sonra, W_n 'de tanımlı $v_i (i = 1, 2, \dots, n)$ orthogonal şebekesi yardımıyla iki invariyan tanımlanmıştır.

Anahtar Sözcükler: Weyl uzayı, Eğrilerin şebekesi, Prolonged kovaryand türev.

* e-posta: nkofoglu@fened.msu.edu.tr, tel: (0212) 236 69 36 / 174

1. INTRODUCTION

An n dimensional manifold W_n is said to be Weyl space if it has a symmetric conformal metric tensor g_{ij} and a symmetric connection ∇_k satisfying the compatibility condition given by the equation

$$\nabla_k g_{ij} = 2T_k g_{ij} \tag{1.1}$$

where T_k denotes a covariant vector field. The symmetric tensor g_{ij} is normalization of the form

$$g_{ij}^v = \lambda^2 g_{ij} \tag{1.2}$$

and the covariant vector field T_k is transformed by the rule

$$T_k^v = T_k + \partial_k \ln \lambda \tag{1.3}$$

where λ is a point function on W_n [1].

Let the coordinates of a Weyl hypersurface of n dimensions and its enveloping space of $(n + 1)$ dimensions be defined by u^k and x^a , respectively and let

$$\partial_k = \frac{\partial}{\partial u^k} . \tag{1.4}$$

A quantity A is called satellite with weight of k of the fundamental tensor g_{ij} .

Under the renormalization of g_{ij} of the form $g_{ij}^v = \lambda^2 g_{ij}$, the quantity A changes according the rule

$$A^* = \lambda^k A . \tag{1.5}$$

The prolonged covariant derivative and the prolonged derivative of the satellite A are defined as follows, respectively

$$\dot{\nabla}_i A = \nabla_i A - k T_i A \tag{1.6}$$

[2] and

$$\dot{\partial}_i A = \partial_i A - k T_i A \tag{1.7}$$

[3] where $\nabla_i A$ is usual derivative of the satellite A and $\partial_i A$ is partial derivative of the satellite A .

Let w be any vector belonging to W_n . If w^i are the contravariant components of the vector w , the covariant derivative of w^i with respect to u^k is

Some Properties Concerning the Hypersurfaces ...

$$\nabla_k \omega^i = \frac{\partial \omega^i}{\partial u^k} + W_{pk}^i \omega^p \quad (1.8)$$

where W_{pk}^i are the coefficients of the connection ∇_k and are defined by the form

$$W_{jk}^i = \left\{ \begin{matrix} i \\ jk \end{matrix} \right\} - (T_k \delta_j^i + T_j \delta_k^i - T_m g_{jk} g^{mi}) \quad (1.9)$$

and $\left\{ \begin{matrix} i \\ jk \end{matrix} \right\}$ are called the second kind Christoffel symbols and are defined as follows

$$\left\{ \begin{matrix} i \\ jk \end{matrix} \right\} = \frac{1}{2} g^{ir} (g_{jr,k} + g_{kr,j} - g_{jk,r}). \quad (1.10)$$

Here, the $g_{jr,k}$ is the partial derivative of g_{jr} with respect to u^k .

A Weyl space is shortly denoted by $W_n(W_{jk}^i, g_{ij}, T_k)$.

Let $W_n(g_{ij}, T_k)$ be a hypersurface of the Weyl space $W_{n+1}(g_{ab}, T_c)$ and x^a ($a = 1, 2, \dots, n+1$) and u^i ($i = 1, 2, \dots, n$) be the coordinates of $W_{n+1}(g_{ab}, T_c)$ and $W_n(g_{ij}, T_k)$, respectively. The metrics of $W_n(g_{ij}, T_k)$ and $W_{n+1}(g_{ab}, T_c)$ are connected by the relations

$$g_{ij} = g_{ab} x_i^a x_j^b \quad (j = 1, 2, \dots, n; b = 1, 2, \dots, n+1) \quad (1.11)$$

where x_i^a is the covariant derivative of x^a with respect to u^i .

The prolonged covariant derivative of A with respect to u^k and x^c are $\dot{\nabla}_k A$ and $\dot{\nabla}_c A$, respectively. These are related by the conditions

$$\dot{\nabla}_k A = x_k^c \dot{\nabla}_c A \quad (k = 1, 2, \dots, n; c = 1, 2, \dots, n+1). \quad (1.12)$$

Let the normal vector field n^a of $W_n(g_{ij}, T_k)$ be normalized by the condition $g_{ab} n^a n^b = 1$. Since the weight of x_i^a is $\{0\}$, the prolonged covariant derivative of x_i^a , relative to u^k , is given by [1]

$$\dot{\nabla}_k x_i^a = \nabla_k x_i^a = \omega_{ik} n^a \quad (1.13)$$

where ω_{ik} are the coefficients of the second fundamental form of $W_n(g_{ij}, T_k)$.

On the other hand, it is easy to see that the prolonged covariant derivative of n^a is given by

$$\dot{\nabla}_k n^a = -\omega_{kl} g^{il} x_i^a. \quad (1.14)$$

By means of (1.13), the prolonged covariant derivative of x_a^j is found to be [4]

$$\dot{\nabla}_k x_a^j = \Omega_k^j n_a. \tag{1.15}$$

Let v_r^a and v_r^i be the contravariant components of the vector field v_r relative to $W_{n+1}(g_{ab}, T_c)$ and $W_n(g_{ij}, T_k)$, respectively. Denoting the components of v^r relative to $W_{n+1}(g_{ab}, T_c)$ and $W_n(g_{ij}, T_k)$ by v_a^r and v_i^r we have [4,5]

$$v_r^a = x_i^a v_r^i, v_a^r = x_a^i v_i^r. \tag{1.16}$$

The prolonged covariant derivatives of the vector field v_r and its reciprocal v^r are, respectively, given by [5]

$$\dot{\nabla}_k v_r^i = T_{r k}^s v_s^i, \dot{\nabla}_k v_i^r = -T_{s k}^r v_i^s. \tag{1.17}$$

2. THE FORMULAS BELONGING TO THE ORTHOGONAL NET

The prolonged covariant derivative of n^a in the direction of v_1^k can be written in the form [6,7]

$$v_1^k \dot{\nabla}_k n^a = k_1^* v_1^a, k_1^* = g_{ab} \left(v_1^k \dot{\nabla}_k n^a \right) v_1^b \tag{2.1}$$

in which

$$g_{ab} n^a v_1^b = 0, g_{ab} n^a n^b = 1, g_{ab} v_1^a v_1^b = 1. \tag{2.2}$$

We call v_1^* the first tangent vector field and $k_1^* v_1^*$ the first curvature vector field and k_1^* the first curvature of C.

Since

$$g_{ab} \left(v_1^k \dot{\nabla}_k n^a \right) n^b = k_1^* g_{ab} v_1^a n^b = 0, \tag{2.3}$$

n^a is perpendicular to $v_1^k \dot{\nabla}_k n^a$.

Taking the prolonged covariant derivative of v_1^* in the direction of v_1^* , we find that

$$v_1^k \dot{\nabla}_k v_1^a = \alpha n^a + \beta v_2^a \tag{2.4}$$

where

Some Properties Concerning the Hypersurfaces ...

$$\mathcal{G}_{ab} n^a v_2^b = 0, \mathcal{G}_{ab} v_1^a v_2^b = 0, \mathcal{G}_{ab} v_2^a v_2^b = 1. \quad (2.5)$$

By using (2.2), (2.3) and (2.5) from (2.4), we obtain $\alpha = -k_1$. Putting $\beta = k_2$, (2.4) becomes

$$v_1^k \dot{\nabla}_k v_1^a = -k_1 n^a + k_2 v_2^a. \quad (2.6)$$

We call v_2 the second tangent vector field and $k_2 v_2$ the second curvature vector field and k_2 the second curvature of C.

Since v_s is perpendicular to v_{s-1} , the equality $\mathcal{G}_{ab} v_s^a v_{s-1}^b = 0$ is satisfied. If we take the prolonged covariant derivative of this equality in the direction of v_1 , we get

$$v_1^k \dot{\nabla}_k v_s^a = -k_s v_{s-1}^a + k_{s+1} v_{s+1}^a \quad (2.7)$$

We call v_{s+1} the (s+1) th. tangent vector field and $k_{s+1} v_{s+1}$ the (s+1) th. curvature vector field and k_{s+1} the (s+1) th. curvature of C.

Proceeding the same way, we finally obtain the derivative of v_n in the form

$$v_1^k \dot{\nabla}_k v_n^a = -k_n v_{n-1}^a \quad (2.8)$$

Generally, the expression (2.8) can be written as

$$v_1^k \dot{\nabla}_k v_p^a = k_{p+1} v_{p+1}^a - k_p v_{p-1}^a \quad (2.9)$$

where $k_{n+1} = 0$ and $k_0 = 0$.

We call v_n the nth. tangent vector field and $k_n v_{n+1}$ the nth. curvature vector field and k_n the nth. curvature of C.

In this way, we obtain n mutually orthogonal vectors v_1, v_2, \dots, v_n at a point P of C which satisfy the condition

$$\mathcal{G}_{ab} v_r^a v_s^b = \delta_r^s \quad (r, s = 1, 2, \dots, n) \quad (2.10)$$

Now, we shall obtain some important results by the first and second order prolonged covariant derivatives of the vector fields of the ennuple:

From (2.7), we have

$$v^l \dot{\nabla}_l v_s^a = -k_s v_{s-1}^a + k_{s+1} v_{s+1}^a$$

If we take the prolonged covariant derivative of both sides of the above equality in the direction of v_1 , we have

$$\begin{aligned} v_s^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) &= v_1^m \left(\dot{\nabla}_m v_1^l \right) \left(\dot{\nabla}_l v_s^a \right) + v_1^l v_1^m \dot{\nabla}_m \left(\dot{\nabla}_l v_s^a \right) \\ &= k_{s+1} \left(k_{s+2} v_{s+2}^a - k_{s+1} v_s^a \right) + v_{s+1}^a v_1^m \left(\dot{\nabla}_m k_{s+1} \right) - k_s \left(k_s v_s^a - k_{s-1} v_{s-2}^a \right) \\ &\quad - v_{s-1}^a v_1^m \left(\dot{\nabla}_m k_s \right) \end{aligned} \tag{2.11}$$

If we multiple (2.11) by $g_{ab} v_{s+1}^b$, we get

$$g_{ab} v_{s+1}^b v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) = v_1^m \left(\dot{\nabla}_m k_{s+1} \right) \tag{2.12}$$

If k_{s+1} is constant, the left hand side of (2.12) is zero and if left hand side of (2.12) is zero,

k_{s+1} is constant.

Hence:

Corollary 2.1

The necessary and sufficient condition that the second order prolonged covariant derivative of the S th. tangent vector field in the direction of v_1 be orthogonal to v_{s+1} is that k_{s+1} be constant.

Let us multiple (2.11) by $g_{ab} v_s^b$. Then we have

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) \right) v_s^b = - \left(k_{s+1}^2 + k_s^2 \right). \tag{2.13}$$

If $k_{s+1} = k_s = 0$, the right hand side of (2.13) is zero. That is, the left hand side of (2.13) is zero. If left hand side of (2.13) is zero, $k_s = k_{s+1} = 0$. From this it follows that

Corollary 2.2

The necessary and sufficient condition that the second order prolonged covariant derivative of the S th. tangent vector field be orthogonal to itself is that both $k_s = 0$ and $k_{s+1} = 0$.

Some Properties Concerning the Hypersurfaces ...

If we multiple (2.11) by $\mathcal{G}_{ab} v_{s-1}^b$, we obtain

$$\mathcal{G}_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) \right) v_{s-1}^b = -v_1^m \dot{\nabla}_m k_s. \quad (2.14)$$

If k_s is constant, the right hand side of (2.14) is zero and conversely, if the left hand side of (2.14) is zero, then k_s is constant.

Hence:

Corollary 2.3

The necessary and sufficient condition that the second order prolonged covariant derivative of the S th. tangent vector field in the direction of v_1 be orthogonal to the $(s-1)$ th.

tangent vector field is that k_s be constant.

If we replace S by $s-1$ in (2.12), we get

$$\mathcal{G}_{ab} v_s^b v_1^m \left(\dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_{s-1}^a \right) \right) = v_1^m \dot{\nabla}_m k_s. \quad (2.15)$$

From (2.14) and (2.15), we obtain

$$\mathcal{G}_{ab} v_s^a v_1^m \left(\dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_{s-1}^b \right) \right) = -\mathcal{G}_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) \right) v_{s-1}^b. \quad (2.16)$$

Therefore:

Corollary 2.4

The components of the second order prolonged covariant derivative along the $(s-1)$ th. tangent vector field is equal to the negative of the components of the $(s-1)$ th. tangent vector field along the S th. tangent vector field.

Since the vector field v_{s-1} is perpendicular to v_s , $\mathcal{G}_{ab} v_{s-1}^a v_s^b = 0$ holds. If we take the prolonged covariant derivative of this equality, we find

$$\mathcal{G}_{ab} v_{s-1}^a \left(v_1^l \dot{\nabla}_l v_s^b \right) + \mathcal{G}_{ab} \left(v_1^l \dot{\nabla}_l v_{s-1}^a \right) v_s^b = 0. \quad (2.17)$$

Again if we take the prolonged covariant derivative of the same equality, we have

$$\begin{aligned} & \mathcal{G}_{ab} v_{s-1}^a \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^b \right) \right) + 2\mathcal{G}_{ab} \left(v_1^m \dot{\nabla}_m v_{s-1}^a \right) \left(v_1^l \dot{\nabla}_l v_s^b \right) + \\ & \mathcal{G}_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_{s-1}^a \right) \right) v_s^b = 0. \end{aligned} \quad (2.18)$$

With the help of (2.17), we have

$$g_{ab} \left(v_1^m \dot{\nabla}_m v_{s-1}^a \right) \left(v_1^l \dot{\nabla}_l v_s^b \right) = 0 \tag{2.19}$$

From here:

Corollary 2.5

The prolonged covariant derivatives of the consecutive tangent vector fields in the direction of v_1 are orthogonal.

Now, if we multiple (2.11) by $g_{ab} v_{s-2}^b$, we obtain

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) \right) v_{s-2}^b = k_s k_{s-1}. \tag{2.20}$$

If k_s or k_{s-1} is zero then the left hand side of (2.20) is zero and if the left hand side of (2.20) is zero, either $k_s = 0$ or $k_{s-1} = 0$.

Hence:

Corollary 2.6

The necessary and sufficient condition that the second order prolonged covariant derivative of the S th. tangent vector field be orthogonal to $(s - 2)$ th. tangent vector field is that either $k_s = 0$ or $k_{s-1} = 0$.

If we multiple (2.11) by $g_{ab} v_{s+2}^b$, we have

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_s^a \right) \right) v_{s+2}^b = k_{s+1} k_{s+2}. \tag{2.21}$$

If $k_{s+1} = 0$ or $k_{s+2} = 0$, then the left hand side of (2.21) is zero. Conversely, if the left hand side of (2.21) is zero, then either $k_{s+1} = 0$ or $k_{s+2} = 0$.

Therefore:

The necessary and sufficient condition that the second order prolonged covariant derivative of the S th. tangent vector field in the direction of v_1 be orthogonal to $(s + 2)$ th. tangent vector field is that either $k_{s+1} = 0$ or $k_{s+2} = 0$.

If S is replaced by $S + 2$ in (2.20), we have

$$g_{ab} v_1^m \left(\dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_{s+2}^a \right) \right) v_s^b = k_{s+1} k_{s+2}. \tag{2.22}$$

If we compare (2.21) and (2.22), we get

Some Properties Concerning the Hypersurfaces ...

$$\mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_s^a \right) \right)_{s+2}^* = \mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^a \right) \right)_s^*. \quad (2.23)$$

We know that $\mathcal{G}_{ab} \mathcal{V}_s^a \mathcal{V}_{s+2}^b = 0$. If we take the prolonged covariant derivative of both

sides equation in the direction of \mathcal{V}_1 , we have $\mathcal{G}_{ab} \mathcal{V}_s^a \mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^b + \mathcal{G}_{ab} \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_s^a \right)_{s+2}^* \mathcal{V}_s^b = 0$.

Again if we take the prolonged covariant derivative of the last equality, we get

$$\begin{aligned} & \mathcal{G}_{ab} \mathcal{V}_s^a \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^b \right) \right) + \mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \mathcal{V}_s^a \right) \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^b \right) + \\ & \mathcal{G}_{ab} \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_s^a \right) \left(\mathcal{V}_1^m \dot{\nabla}_m \mathcal{V}_{s+2}^b \right) + \mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_s^a \right) \right)_{s+2}^* \mathcal{V}_s^b = 0. \end{aligned} \quad (2.24)$$

Using (2.23) and (2.24), we obtain

$$\begin{aligned} & \mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \mathcal{V}_s^a \right) \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^b \right) = -\mathcal{G}_{ab} \mathcal{V}_s^a \mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_{s+2}^b \right) = \\ & -\mathcal{G}_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_s^a \right) \right)_{s+2}^* \mathcal{V}_s^b \end{aligned} \quad (2.25)$$

From here:

Corollary 2.7

The necessary and sufficient conditions that the prolonged covariant derivative of \mathcal{V} in the direction of \mathcal{V}_1 be orthogonal to the prolonged covariant derivative of \mathcal{V} is that either the second order prolonged covariant derivative of \mathcal{V} be orthogonal to \mathcal{V} or the second order prolonged covariant derivative of \mathcal{V} be orthogonal to \mathcal{V} .

Now, we consider a particular case. For example; the case of $S = 1$. Then we obtain from (2.11) that

$$\begin{aligned} & \mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_1^a \right) = k_2 \left(k_3 \mathcal{V}_3^a - k_2 \mathcal{V}_1^a \right) + \mathcal{V}_2^a \mathcal{V}_1^m \left(\dot{\nabla}_m k_2 \right) - \\ & k_1 \left(k_1 \mathcal{V}_1^a \right) - n^a \mathcal{V}_1^m \dot{\nabla}_m k_1 \end{aligned} \quad (2.26)$$

Multiplying (2.26) by $g_{ab} v_3^b$, the following equality is found

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_1^a \right) \right) v_3^b = k_2 k_3. \tag{2.27}$$

If $k_2 = 0$ or $k_3 = 0$, then the left hand side of (2.40) is zero. Conversely, if the left hand side of (2.27) is zero then either $k_2 = 0$ or $k_3 = 0$.

Therefore:

Corollary 2.8

If $k_2 = 0$ or $k_3 = 0$, the second order prolonged covariant derivative of the first tangent vector field in the direction of v_1 is orthogonal to the third tangent vector field. The converse is also true.

If we multiple (2.26) by $g_{ab} v_1^b$, we get

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_1^a \right) \right) v_1^b = - (k_2^2 + k_1^2). \tag{2.28}$$

If $k_2 = 0 = k_1$, then the left hand side of (2.28) is zero. Conversely, if the left hand side of (2.28) is zero, then $k_2 = 0 = k_1$.

Hence:

Corollary 2.9

$k_2 = 0 = k_1$ is the necessary and sufficient condition for the orthogonality of the second order prolonged covariant derivative of the first tangent vector field in the direction of v_1 to itself.

If we multiple (2.26) by $g_{ab} v_1^b$, we get

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_1^a \right) \right) v_2^b = v_1^m \dot{\nabla}_m k_2. \tag{2.29}$$

If $k_2 = \text{constant}$, then the left hand side of (2.29) is zero. Conversely, if the left hand side of (2.29) is zero, we obtain $k_2 = \text{constant}$.

From here:

Corollary 2.10

The necessary and sufficient condition that the second order prolonged covariant derivative of the first tangent vector field in the direction of v_1 be orthogonal to the second tangent vector field is that k_2 be constant.

Some Properties Concerning the Hypersurfaces ...

Multiplying (2.26) by $g_{ab} n^b$, we have

$$g_{ab} \left(v_1^m \dot{\nabla}_m \left(v_1^l \dot{\nabla}_l v_1^a \right) \right) n^b = -v_1^m \dot{\nabla}_m k_1. \quad (2.30)$$

If k_1 is constant, then the left hand side of (2.30) is zero. Conversely, if the left hand side of (2.30) is zero, then k_1 is constant.

Therefore:

Corollary 2.11

$k_1 = \text{constant}$ is the necessary and sufficient condition that the second order prolonged covariant derivative of the first tangent vector field in the direction of v_1 be orthogonal to the normal vector field.

Hence:

If the right hand side of (2.28) is zero, then the right hand side of (2.27), (2.29) and (2.30) are also zero.

From this it follows

Corollary 2.12

If the second order prolonged covariant derivative of the first tangent vector field in the direction of v_1 is orthogonal to itself, then it is also orthogonal to the normal vector field, the second tangent vector field and the third tangent vector field.

Multiplying (2.6) by $g_{ab} n^b$, we get

$$g_{ab} \left(v_1^l \dot{\nabla}_l v_1^a \right) n^b = -k_1 \quad (2.31)$$

If $k_1 = 0$, then the left hand side of (2.31) is zero and the converse of it is also true.

Hence:

Corollary 2.13

The necessary and sufficient condition that the prolonged covariant derivative of the first tangent vector field in the direction of v_1 be orthogonal to the normal vector field is that k_1 be zero.

If we multiple (2.6) by $g_{ab} v_2^b$, we obtain

$$g_{ab} \left(v_1^l \dot{\nabla}_l v_1^a \right) v_2^b = k_2. \quad (2.32)$$

If $k_2 = 0$, then the left hand side of (2.32) is zero. The converse is also true. Thus, we have

Corollary 2.14

The necessary and sufficient condition that the first order prolonged covariant derivative of the first tangent vector field in the direction of \mathcal{V}_1 be orthogonal to the second tangent vector field is that k_2 be zero.

If $k_1 = 0$ and $k_2 = 0$, then the right hand side of (2.31) and (2.32) are zero. In this case, the right hand side of (2.28) is zero.

Therefore, we have

Corollary 2.15

If the second order prolonged covariant derivative of the first tangent vector field is orthogonal to itself, then the first order prolonged covariant derivative of the first tangent vector field is orthogonal to the normal vector field as well as to the second tangent vector field.

We have seen from (2.32) that if $k_2 = 0$ the first order prolonged covariant derivative of the first tangent vector field is orthogonal to the second tangent vector field. The converse of it is also true. But from (2.27) for $k_2 = 0$, it is seen that the second order prolonged covariant derivative of the first tangent vector field is orthogonal to the third tangent vector field. Hence:

Corollary 2.16

If the first order prolonged covariant derivative of the first tangent vector field in the direction of \mathcal{V}_1 is orthogonal to the second tangent vector field, then the second order prolonged covariant derivative of the first tangent vector field in the direction of \mathcal{V}_1 is orthogonal to the third tangent vector field.

We know that the equation of (2.28) had been expressed as

$$g_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_1^a \right) \right) \mathcal{V}_1^b = - (k_2^2 + k_1^2). \tag{2.33}$$

If k_2 is constant, then from (2.29) the second order prolonged covariant derivative of the first tangent vector field in the direction of \mathcal{V}_1 is orthogonal to the second tangent vector field.

If k_1 is constant, then from (2.30) the second order prolonged covariant derivative of the first tangent vector field in the direction of \mathcal{V}_1 is orthogonal to the normal vector field.

If the conditions $k_1 = \text{constant}$ and $k_2 = \text{constant}$ are satisfied, then we obtain from (2.33) that

$$g_{ab} \left(\mathcal{V}_1^m \dot{\nabla}_m \left(\mathcal{V}_1^l \dot{\nabla}_l \mathcal{V}_1^a \right) \right) \mathcal{V}_1^b = \text{constant}.$$

Corollary 2.17

If the second order prolonged covariant derivative of the first tangent vector field is orthogonal both to the normal vector field and to the second tangent vector, then this and the first tangent vector field cut each other under a constant angle.

Some Properties Concerning the Hypersurfaces ...

3. THE DERIVATIVE FORMULAS FOR A GEODESIC TANGENT

Definition 3.1: Let C be a curve in the Wey hypersurface W_n and let \mathcal{V}_1 be the tangent vector field of C . If the prolonged covariant derivative of \mathcal{V}_1 in the direction of itself is zero, then C is called geodesic, i.e.

$$\mathcal{V}_1^k \dot{\nabla}_k \mathcal{V}_1^l = 0.$$

Let us consider the geodesic tangent vector field of the curve C at the point P and let us denote it by C_g . Furthermore, let us denote the tangent vector field, the principal normal vector field and the binormal vector field belonging to C_g by $\bar{\mathcal{V}}, \bar{\mathcal{N}}, \bar{\mathcal{N}}_2$, respectively

We know from (2.1) that

$$\mathcal{V}_1^k \dot{\nabla}_k n^a = k_1^* \mathcal{V}_1^a \quad (3.1)$$

We can write

$$\mathcal{V}_1^k \dot{\nabla}_k n^a = \bar{\mathcal{V}}^k \dot{\nabla}_k n^a = \tau_2 \bar{\mathcal{N}}_2^a - \tau_1 \bar{\mathcal{V}}_1^a \quad (3.2)$$

from Darboux_Ribocour Equations, where τ_1 and τ_2 are the first and the second curvature of the geodesic tangent to the curve C , respectively, τ_1 and τ_2 are the normal curvature and the geodesic torsion of the curve C , respectively.

From (3.1) and (3.2), we get

$$k_1^* \mathcal{V}_1^a = \tau_2 \bar{\mathcal{N}}_2^a - \tau_1 \bar{\mathcal{V}}_1^a. \quad (3.3)$$

If we multiple (3.3) by itself, we obtain

$$k_1^2 = \tau_1^2 + \tau_2^2. \quad (3.4)$$

Therefore, we can state the following theorem:

Theorem: 3.1.

If any two of the following properties for a curve in a hypersurface W_n of Weyl space W_{n+1} are satisfied, then the third also holds:

- i)* The first curvature of the geodesic tangent vanishes.
- ii)* The geodesic torsion of the curve is zero.
- iii)* The first curvature of the curve C is zero

Theorem: 3.2.

If the curve C is an asymptotic line, then the prolonged covariant derivative of the normal vector field in the direction of \mathcal{V}_1 is orthogonal to the curve.

Proof: 3.2.

Let C be an asymptotic line. Then the normal curvature of C is zero, that is, $\kappa_{11} = 0$.

For C_g ,

$$\kappa_{11} = \tau_1 = 0 \tag{3.5}$$

is satisfied.

We know that

$$\kappa_{11} = -g_{ab} \left(v_1^k \dot{\nabla}_k n^a \right) v_1^b. \tag{3.6}$$

From (3.5) and (3.6), we see that $v_1^k \dot{\nabla}_k n^a$ is orthogonal to v_1^* .

The proof is completed.

On the other hand, we know that $v_1^k \dot{\nabla}_k n^a = \tau_2 \bar{n}_2^a - \tau_1 \bar{v}_1^a$ from (3.2). Since

$$g_{ab} \left(v_1^k \dot{\nabla}_k n^a \right) \bar{v}_1^b = 0, \text{ we get}$$

$$\tau_1 = 0. \tag{3.7}$$

This says

$$k_1 = \tau_2. \tag{3.8}$$

Also from (3.3)

$$v_1^a = \bar{n}_2^a \tag{3.9}$$

From (3.8):

Corollary 3.1

The product of the prolonged covariant derivative of the normal vector field along asymptotic line by itself is the geodesic torsion of the asymptotic line.

4. ON THE HYPERSURFACES MEETING UNDER A CONSTANT ANGLE

We consider a curve C with the tangent vector field v_1 which is common to two hypersurfaces

W_n and \bar{W}_n . Let n and \bar{n} be the normal vector fields with respect to these hypersurfaces. If these hypersurfaces meet at a constant angle then the following condition is satisfied:

$$v_1^k \dot{\nabla}_k (g_{ab} n^a \bar{n}^b) = 0 \tag{4.1}$$

From this it follows that

$$g_{ab} \left\{ v_1^k (\dot{\nabla}_k n^a) \right\} \bar{n}^b + g_{ab} n^a \left\{ v_1^k \dot{\nabla}_k \bar{n}^b \right\} = 0. \tag{4.2}$$

Some Properties Concerning the Hypersurfaces ...

Since W_n and \overline{W}_n meet under a constant angle, C is a line of curvature for both hypersurfaces, so that, $v_1^k \dot{\nabla}_k n^a = k_1 v_1^a$ where v_1^a are the components of the tangent vector field to the curve with respect to W_{n+1} and k_1^2 is the inner product of the prolonged covariant derivative of the normal vector field n in the direction of v_1 by itself.

Also $\overline{v}_1^k \dot{\nabla}_k \overline{n}^a = \overline{k}_1 \overline{v}_1^a$ where \overline{v}_1^a are the components of the tangent vector field to the curve with respect to \overline{W}_{n+1} and \overline{k}_1^2 is the inner product of the prolonged covariant derivative of the normal vector field \overline{n} in the direction of \overline{v}_1 by itself.

From the above information and (4.2), we obtain

$$g_{ab} \left(k_1 v_1^a \right) \cdot \overline{n}^b + g_{ab} n^a \left(\overline{k}_1 \overline{v}_1^b \right) = 0,$$

(4.3) or

$$\frac{k_1}{\overline{k}_1} = - \frac{g_{ab} n^a \overline{v}_1^b}{g_{ab} v_1^a \overline{n}^b} = - \frac{\cos \angle \left(n^a, \overline{v}_1^b \right)}{\cos \angle \left(v_1^a, \overline{n}^b \right)}. \quad (4.4)$$

We can express this as:

Theorem : 4.1.

If a curve C is common to two hypersurfaces W_n and \overline{W}_n of the Weyl space $W_{n+1}(g_{ab}, T_c)$ such that they meet under a constant angle along C, then $\frac{k_1}{\overline{k}_1}$ is a gauge invariant for $W_{n+1}(g_{ab}, T_c)$.

5. AN INVARIANT ASSOCIATED WITH AN ORTHOGONAL ENNUPLE IN A WEYL HYPERSURFACE

Theorem: 5.1.

The sum $\sum_{r=1}^n k_r^2$ where k_r^2 is the inner product of the prolonged covariant derivative of the normal vector field to a Weyl hypersurface in the direction, of the rth vector of an orthogonal ennuple by itself, is an invariant.

Proof:

Let us denote the orthogonal ennuple in W_n by (v_1, v_2, \dots, v_n) . The prolonged covariant derivative of the normal vector field in the directions of the vectors of the orthogonal ennuple can be expressed, by (1.14), as

$$v_r^k \dot{\nabla}_k n^a = -v_r^k \omega_{kl} g^{il} x_i^a \quad (r = 1, 2, \dots, n). \tag{5.1}$$

From this we obtain

$$\begin{aligned} \sqrt{g_{ab} (v_r^k \dot{\nabla}_k n^a) (v_r^l \dot{\nabla}_l n^b)} &= \sqrt{g_{ab} (-v_r^k \omega_{kj} g^{ij} x_i^a) (-v_r^l \omega_{lm} g^{ml} x_l^b)} \\ &= \sqrt{g^{mj} v_r^k v_r^l \omega_{kj} \omega_{lm}} \end{aligned} \tag{5.2}$$

with the help of (1.11) and (5.1). Let us denote this scalar by k_r , i.e. $k_r^2 = g^{mj} v_r^k v_r^l \omega_{kj} \omega_{lm}$.

If we take the sum of the squares with respect to r , we find from (5.2) that

$$\sum_{r=1}^n k_r^2 = \sum_{r=1}^n g^{mj} v_r^k v_r^l \omega_{kj} \omega_{lm} = g^{mj} g^{kl} \omega_{kj} \omega_{lm}, \tag{5.3}$$

since $\sum_{r=1}^n v_r^k v_r^l = g^{kl}$ for the vector fields of an orthogonal ennuple.

This shows that $\sum_{r=1}^n k_r^2$ is an invariant. The proof of the theorem is completed.

I am thankful to Prof. Dr. Leyla Zeren Akgün for her guidance.

REFERENCES

- [1] Norden, A. : Affinely Connected Spaces, GRMFL, Moscow, (1976).
- [2] Norden, A. , Yafarov, S. : Theory of Non-geodesic Vector Fields in Two Dimensional Affinely Connected Spaces, Izv. , Vuzov, Math. , No. 12, 29-34, (1974).
- [3] Hlavaty, V. : Les Courbes de la Variete W_n , Memor. Sci. Math. , Paris, (1934).
- [4] Uysal, S. A. , Özdeğer, A. : On the Chyshev Nets in a Hypersurface of a Weyl Space, Journal of Geometry, V.51, 171-177, (1994).
- [5] Tsareva, B. , Zlatanov, G. : On the Geometry of the Nets in the n -Dimensional Space of Weyl, Journal of Geometry, Vol. 38, 182-197, (1955).
- [6] Kaul, R. N. : Formulae Corresponding to Frenet's Formulae, Acad. Roy. Belgique, 41, 1292-1304, (1955).
- [7] Akgün, L. Z. : Frenet Formulas For Curves in a Generalized Weyl Space, Ganita, Vol.51, No.2, 149-164, (2000).